

α -Representation for QCD

Richard Hong Tuan

Laboratoire de Physique Théorique et Hautes Energies¹
Université de Paris XI, Bâtiment 210, F-91405 Orsay Cedex, France

Abstract

An α -parameter representation is derived for gauge field theories. It involves, relative to a scalar field theory, only constants and derivatives with respect to the α -parameters. Simple rules are given to obtain the α -representation for a Feynman graph with an arbitrary number of loops in gauge theories in the Feynman gauge.

LP THE Orsay 98-45
July 1998

¹Laboratoire associé au Centre National de la Recherche Scientifique - URA D0063

An α -parameter representation may be useful for a variety of purposes. It was essential in deriving [1] string motivated one-loop rules for connecting ϕ^3 amplitudes to QCD amplitudes. Starting from a ϕ^3 α -parameter representation one gets a QCD α -parameter representation by multiplying the ϕ^3 expression by a “reduced” factor which is calculated by methods of string-theory. However, going to more than one-loop level appears difficult. Here, we propose an α -representation for gauge theories which can be easily realized by simple rules involving derivatives with respect to the α -parameters and *for any number of loops*. Such a representation may also be useful to compute Regge trajectories for QCD as we have done [2] for ϕ^3 .

Let us start with the amplitude for a Feynman graph G of a scalar field theory with I internal lines and V vertices [3]. We define the incidence matrix $\varepsilon_{v\ell}$ with indices running over vertices v and propagators ℓ as

$$\varepsilon_{v\ell} = \begin{cases} 1 & \text{if } v \text{ is the starting point of } \ell \\ -1 & \text{if } v \text{ is the endpoint of } \ell \\ 0 & \text{if } \ell \text{ is not incident with } v \end{cases} . \quad (1)$$

Let us denote by $P_v = \sum p_{kv}$ the sum of incoming external momenta at the vertex v . Then, in a theory without derivative couplings, i.e. a scalar field theory, we get for the Feynman amplitude F_G of G (in $d = 4$ dimension)

$$F_G = [C(G)/S(G)]$$

$$\int \left[\prod_{\ell=1}^I \frac{d^4 k_\ell}{2\pi^4} \frac{i}{k_\ell^2 - m_\ell^2 + i\varepsilon} \right] \prod_{v=1}^V (2\pi)^4 \delta^4 \left(P_v - \sum_{\ell=1}^I \varepsilon_{v\ell} k_\ell \right) \quad (2)$$

where $C(G)$ contains the vertices factors dependence and $S(G)$ is the symmetry factor. The four-momentum k_ℓ is oriented as the line ℓ is. Now,

$$i/(p^2 - m^2 + i\varepsilon) = \int_0^\infty d\alpha \exp \left[i\alpha(p^2 - m^2 + i\varepsilon) \right] \quad (3)$$

$$(2\pi)^4 \delta^4 \left(P_v - \sum_{\ell=1}^I \varepsilon_{v\ell} k_\ell \right) = \int d^4 y_v \exp \left[-iy_v \cdot \left(P_v - \sum_{\ell} \varepsilon_{v\ell} k_\ell \right) \right] \quad (4)$$

Inserting (3) and (4) in (2) and interchanging integrations order (regularization is needed for divergent integrals) we get for each $d^4 k_\ell$ integration

$$\begin{aligned} & \int d^4 k_\ell / (2\pi)^4 \exp \left[i\alpha_\ell \left(k_\ell^2 + \alpha_\ell^{-1} \sum_i y_v \cdot \varepsilon_{v\ell} k_\ell \right) \right] \\ &= \int d^4 k_\ell / (2\pi)^4 \exp \left\{ i\alpha_\ell \left[\left(k_\ell + (2\alpha_\ell)^{-1} \sum_v y_v \varepsilon_{v\ell} \right)^2 - (4\alpha_\ell^2)^{-1} \left(\sum_v y_v \varepsilon_{v\ell} \right)^2 \right] \right\} . \end{aligned} \quad (5)$$

The k_ℓ dependence is now concentrated in the first term of the exponential. This is crucial for what follows. Derivative couplings manifest themselves through k_ℓ^μ factors. Uneven powers of k_ℓ cancel after making the variable substitution

$$k_\ell \rightarrow k_\ell - (2\alpha_\ell)^{-1} \sum_v y_v \varepsilon_{v\ell} \quad , \quad (6)$$

the exponential is then simplified and any k_ℓ^2 factor with transform, giving a k_ℓ -dependent integral

$$\int d^4 k_\ell / (2\pi)^4 \exp \left(i\alpha_\ell k_\ell^2 \right) \left[k_\ell - (2\alpha_\ell)^{-1} \sum_v y_v \varepsilon_{v\ell} \right]^2 . \quad (7)$$

However, this is precisely the kind of factor QCD will exhibit. If, in Feynman gauge,

$$\begin{aligned} \mathcal{L}_0 = & -1/4 \left(\partial_\mu A_\nu^a - \partial_\nu A_\mu^a \right) \left(\partial^\mu A^{a\nu} - \partial^\nu A^{a\mu} \right) \\ & -1/2 \left(\partial^\mu A_\mu^a \right)^2 + \left(\partial^\mu \chi^{a*} \right) \left(\partial_\mu \chi^a \right) \\ & + \bar{\psi} \left(i\gamma^\mu \partial_\mu - m \right) \psi \end{aligned} \quad (8.a)$$

$$\begin{aligned} \mathcal{L}_1 = & -g/2 f^{abc} \left(\partial_\mu A_\nu^a - \partial_\nu A_\mu^a \right) A^{b\mu} A^{c\nu} \\ & -g^2/4 f^{abc} f^{cde} A_\mu^a A_\nu^b A^{c\mu} A^{d\nu} \\ & -g f^{abc} \left(\partial^\mu \chi^{a*} \right) \chi^b A_\mu^c + g \bar{\psi} T^a \gamma^\mu \psi A_\mu^a \end{aligned} \quad (8.b)$$

are respectively the free and the interacting part of the QCD Lagrangian, one remembers that ∂_μ is contracted with $A^{b\mu}$, a field upon which ∂_μ does not act in (8.b). If there is another ∂_μ acting on the same propagator, but at the other end, a dot-product $\partial_\mu \partial^\mu$ is made (contracting indices along propagators joining the ends of the first propagator on which the considered ∂_μ 's act) and we get, Fourier transforming (putting aside coupling constants)

$$(-ik_{\mu\ell}) (ik_\ell^\mu) = k_\ell^2 . \quad (9)$$

Making the variable change we therefore get (7) or, eliminating odd powers of k_ℓ which give vanishing contributions,

$$\int d^4 k_\ell / (2\pi)^4 \exp(i\alpha_\ell k_\ell^2) \left[k_\ell^2 + (2\alpha_\ell)^{-2} \left(\sum_v y_v \varepsilon_{v\ell} \right)^2 \right] . \quad (10)$$

Now, the crucial remark is that the bracket in (10) can be obtained *by the derivation with respect to $i\alpha_\ell$ of the integrand of (5)*. This is a very simple rule indeed to obtain the kinematical factor associated with a derivative coupling : *excluding the factor $\exp(-i\alpha_\ell m_\ell^2)$, we simply have to take the derivative of the integrand with respect to $i\alpha_\ell$* . Of course, there can be other dot-products arising involving k_μ 's like $(k_\mu \varepsilon^\mu)(k_\nu \varepsilon^\nu)$, $(k_{1\mu} k_2^\mu)(k_{2\nu} k_1^\nu)$. However, because, in d dimensions,

$$\int d^d k_\ell k_\ell^\mu k_\ell^\nu f(k_\ell^2) = (g^{\mu\nu}/d) \int d^d k_\ell k_\ell^2 f(k_\ell^2) \quad (11)$$

all dot-products will give factors proportional to k_ℓ^2 and the rule given above will hold up to a d -dependent coefficient. For instance, for $(k_1 \cdot k_2)(k_2 \cdot k_1)$ this coefficient will be $d/d^2 = d^{-1}$ in factor of the operator $\partial/\partial(i\alpha_1) \partial/\partial(i\alpha_2)$.

If we perform the $\int d^d k_\ell$ integrals (no derivation being done) we get, in d dimensions (in order to allow for dimensional regularization)

$$F_G = [C(G)/S(G)] \int \prod_{v=1}^V d^d y_v \int_0^\infty \left[\prod_{\ell=1}^I d\alpha_\ell i^{-(d-2)/2} (4\pi\alpha_\ell)^{-d/2} \right. \\ \left. \exp \left\{ -i \left[\alpha_\ell m_\ell^2 + \left(\sum_{v=1}^V y_v \varepsilon_{v\ell} \right)^2 / (4\alpha_\ell) \right] \right\} \right] \exp \left(-i \sum_{v=1}^V y_v \cdot P_v \right) . \quad (12)$$

Making the change of variable

$$\begin{aligned} z_i &= y_i - z_v \quad i = 1, \dots, V-1 \\ z_V &= y_V \end{aligned} \tag{13}$$

because $\sum_{v=1}^V \varepsilon_{v\ell} = 0$ (ℓ connects two and only two vertices), z_V only appears in the last exponent of (12) and the integration over z_V yields a factor $\delta^d\left(\sum_v P_v\right)$ implying energy-momentum conservation. There remains $V-1$ independent variables z_v to be integrated over. Separating the z_v dependent factors, the z_v integral reads

$$\begin{aligned} I_H &= \int \left[\prod_{v=1}^{V-1} d^d z_v \right] H(\{j_\ell\}) \\ &\exp \left\{ -i \left[\sum_{\ell} (4\alpha_\ell)^{-1} \left(\sum_{v=1}^{V-1} z_v \varepsilon_{v\ell} \right)^2 + \sum_{v=1}^{V-1} z_v \cdot P_v \right] \right\} \end{aligned} \tag{14}$$

where $H(\{j_\ell\})$ represents factors coming from the derivative couplings through j_ℓ 's defined as

$$j_\ell = (2\alpha_\ell)^{-1} \sum_{v=1}^{V-1} z_v \varepsilon_{v\ell} \quad . \tag{15}$$

As previously quoted, to k_ℓ^2 is associated $(k_\ell - j_\ell)^2$ through the change of variable (6), giving $k_\ell^2 + j_\ell^2$, eliminating the term linear in k_ℓ . The j_ℓ^2 term is obtained by letting $\partial/\partial(i\alpha_\ell)$ act upon (14). However, we may have other sources of j_ℓ dependence occurring in $H(\{j_\ell\})$. Let us explain this.

If G were a tree, then G would have only $V - 1$ propagators and a variable change (with Jacobian equal to one)

$$z_v \rightarrow \sum_{v=1}^{V-1} z_v \varepsilon_{v\ell} \quad (16)$$

would be feasible while keeping j_ℓ 's independent of each other as far z_v dependence is concerned. Then, any factor $j_\ell \cdot j_{\ell'}$, coming for instance from a factor $(k_\ell - j_\ell) \cdot (k_{\ell'} - j_{\ell'})$, with $\ell \neq \ell'$, would factorize into independent integrals, making the evaluation of (14) straightforward. But, as soon as G contains loops, because around a loop $\mathcal{L}$

$$\sum_{\ell \in \mathcal{L}} \varepsilon_\ell \left(\sum_{v=1}^{V-1} z_v \varepsilon_{v\ell} \right) = 0 \quad , \quad \varepsilon_\ell = \pm 1 \quad (17)$$

if ℓ and ℓ' belong to the same loop $\mathcal{L}$, j_ℓ and $j_{\ell'}$ are not independent and we have a more difficult integration to do. We now treat the general case where loops are present. We first eliminate the linear term in the exponential of (11) through a variable change

$$z_v \rightarrow z_v - \beta_v \quad (18)$$

which yields $V - 1$ equations

$$-(1/2) \sum_{v'=1}^{V-1} \beta_{v'} \sum_{\ell} \varepsilon_{v'\ell} \alpha_\ell^{-1} \varepsilon_{v\ell} + P_v = 0 \quad . \quad (19)$$

The $V - 1$ by $V - 1$ symmetric matrix d_G defined as

$$[d_G(\alpha)]_{v_1 v_2} = \sum_{\ell} \varepsilon_{v_1 \ell} \alpha_{\ell}^{-1} \varepsilon_{v_2 \ell} \quad (20)$$

is known to be non-singular and its determinant reads

$$\Delta_G(\alpha) = \sum_{\mathcal{T}} \prod_{\ell \in \mathcal{T}} \alpha_{\ell}^{-1} \quad (21)$$

where the sum runs over all spanning trees $\mathcal{T}$ of G (a spanning tree is a tree incident with every vertex of G). We therefore get

$$\beta_v = 2 \sum_{v'=1}^{V-1} P_{v'} [d_G(\alpha)]_{v'v}^{-1} \quad (22)$$

which means that a factor $j_{\ell} \cdot j_{\ell'}$ is transformed according to

$$j_{\ell} \cdot j_{\ell'} \rightarrow j_{\ell} \cdot j_{\ell'} + (4\alpha_{\ell} \alpha_{\ell'})^{-1} \left(\sum_{v=1}^{V-1} \beta_v \varepsilon_{v\ell} \right) \left(\sum_{v'=1}^{V-1} \beta_{v'} \varepsilon_{v'\ell'} \right) \quad (23)$$

+terms linear in j_{ℓ} or $j_{\ell'}$.

We now make an orthogonal transformation in order to diagonalize $d_G(\alpha)$. Then, performing the integration over the $V-1$ z_v vectors, the terms linear in j_{ℓ} and $j_{\ell'}$ vanish and for $H(\{j_{\ell}\}) = j_{\ell} \cdot j_{\ell'}$ we get (two derivations of (14) with respect to iP_v also do the job)

$$I_H = I_0 (4\alpha_{\ell} \alpha_{\ell'})^{-1} \left\{ -2id \sum_{v_1, v_2=1}^{V-1} \varepsilon_{v_1 \ell} [d_G(\alpha)]_{v_1 v_2}^{-1} \varepsilon_{v_2 \ell'} \right. \\ \left. + \left(\sum_{v_1=1}^{V-1} \beta_{v_1} \varepsilon_{v_1 \ell} \right) \cdot \left(\sum_{v_2=1}^{V-1} \beta_{v_2} \varepsilon_{v_2 \ell'} \right) \right\} \quad (24)$$

with

$$I_0 = i^{(V-1)(d-2)/2} (4\pi)^{(V-1)d/2} [\Delta_G(\alpha)]^{-d/2} \exp \left\{ i \sum_{v_1, v_2=1}^{V-1} P_{v_1} [d_G(\alpha)]_{v_1 v_2}^{-1} \cdot P_{v_2} \right\} \quad (25)$$

I_0 being I_H for $H(\{j_\ell\}) = 1$. One remarks that, of course, if ℓ and ℓ' belong to the same spanning tree, I_H reduces to its second term in (24). Now, if ℓ and ℓ' belong to the same loop $\mathcal{L}$, we get according to (17),

$$j_\ell \cdot j_{\ell'} = - (4\alpha_\ell \alpha_{\ell'})^{-1} \left(\sum_{v_1=1}^{V-1} z_{v_1} \varepsilon_{v_1 \ell} \right) \varepsilon_{\ell'} \sum_{\substack{\ell_1 \neq \ell' \\ \ell_1 \in \mathcal{L}}} \varepsilon_{\ell_1} \left(\sum_{v_2=1}^{V-1} z_{v_2} \varepsilon_{v_2 \ell_1} \right) \quad . \quad (26)$$

Cutting $\mathcal{L}$ at ℓ' , we get that all $\ell_1 \neq \ell'$ belonging to $\mathcal{L}$ are on a spanning tree of G . Then, the crossed terms in (26) with $\ell_1 \neq \ell$ will not contribute, leaving us only with

$$-\varepsilon_\ell \varepsilon_{\ell'} (4\alpha_\ell \alpha_{\ell'})^{-1} \left(\sum_{v=1}^{V-1} z_v \varepsilon_{v \ell} \right)^2 \quad (27)$$

because all j_ℓ 's on a spanning tree are independent. As (27) is proportional to j_ℓ^2 it can be obtained by a derivation and (24) will be written (using (14))

$$I_H = -\varepsilon_\ell \varepsilon_{\ell'} (\alpha_\ell / \alpha_{\ell'}) \partial / \partial (i\alpha_\ell) I_0 \quad (28)$$

as can be easily checked against (24) by a direct evaluation where we remark that the first term in (24) comes from derivating $[\Delta_G(\alpha)]^{-d/2}$ and the second

one from derivating the exponential. Now, instead of dot-products one may have tensor products for j_ℓ 's coming for instance from the dot-products $(k_1 - j_1) \cdot (k_2 - j_2) (k_2 - j_2) \cdot (k_1 - j_1)$. However, after the variable-shift (18) has been made the formula (11) can be applied to j_ℓ 's as well as to k_ℓ 's. We can now give the procedure for writing, in the Feynman gauge, the α -parameter expression for any QCD graph amplitude :

- i) Write the Feynman numerator which is a polynomial in dot-products and tensor-products of P_v 's, k_ℓ 's and $g^{\mu\nu}$'s.
- ii) Replace each internal momentum k_ℓ by $k_\ell - j_\ell$ in the Feynman numerator.
- iii) Choose a spanning tree $\mathcal{T}$ on G , for which the j_ℓ 's will be independent.
- iv) Replace each j_ℓ of a propagator not belonging to $\mathcal{T}$ by its expression as a function of j_ℓ 's belonging to $\mathcal{T}$ (use relation (17) for that purpose).
- v) For each monomial of the Feynman numerator the product of j_ℓ 's and k_ℓ 's should be replaced by an operator product $\hat{\mathcal{O}}$ according to the following rules (n is a positive integer)

$$k_\ell \rightarrow 0 \tag{29.a}$$

$$k_\ell \otimes k_\ell \rightarrow g^{\mu\nu}/d \quad \partial/\partial(i\alpha_\ell) \quad \text{acting on} \quad \left[\prod_{\ell=1}^I \alpha_\ell \right]^{-d/2} \tag{29.b}$$

$$(j_\ell \otimes j_\ell)^n \rightarrow \left[g^{\mu\nu}/d \quad \partial/\partial(i\alpha_\ell) + b_\ell^\mu b_\ell^\nu - (g^{\mu\nu}/d) b_\ell^2 \right]^n \quad \text{acting on} \quad I_0 \tag{29.c}$$

$$(j_\ell)^{2n+1} \rightarrow \sum_{p=0}^n C_{2n+1}^{2(n-p)} (-b_\ell)^{2(n-p)+1} (j_\ell \otimes j_\ell)^p \text{ acting on } I_0 \quad (29.d)$$

where $b_\ell = (2\alpha_\ell)^{-1} \sum_{v=1}^{V-1} \beta_v \varepsilon_{v\ell}$. In (29.d) $(j_\ell)^{2n+1}$ is a tensor product. This gives the following expression for F_G^{QCD} in d dimensions

$$F_G^{QCD} = [C(G)/S(G)] i^L (4\pi i)^{-Ld/2} \int_0^\infty \left[\prod_{\ell=1}^I d\alpha_\ell \exp(i\alpha_\ell m_\ell^2) \right] \\ \sum_{\{\hat{\mathcal{O}}\}} \hat{\mathcal{O}} \left[\prod_{\ell=1}^I \alpha_\ell \right]^{d/2} [\Delta_G(\alpha)]^{-d/2} \exp \left\{ i \sum_{v_1, v_2}^{V-1} P_{v_1} [d_G(\alpha)]_{v_1 v_2}^{-1} \cdot P_{v_2} \right\} \quad (30)$$

where m_ℓ is infinitesimal for vector particles, i.e. gluons and $\{\hat{\mathcal{O}}\}$ is the ensemble of monomials in the Feynman numerator. In (29.c) and (29.d) n is at most equal to two because at most three loops can share the same propagator.

We have used [4] the representation (30) in the calculations of the one-loop gluon self-energy and obtained the same result as in the conventional approach.

References

- [1] Z. Bern and D. C. Dunbar, Nucl. Phys. **B379** (1992) 562.
- [2] R. Hong Tuan, Phys. Lett. **B393** (1997) 84 ; preprint LPTHE 97/38, Orsay (1997), hep-ph 9712460.
- [3] C. Itzykson and J. B. Zuber, Quantum Field Theory, McGraw-Hill, p. 294.
- [4] R. Hong Tuan, to be published.